

WATER

Written & Compiled by:

Jim Lafley, DCR Division of Water Supply Protection, Education Coordinator

Jim Lagacy, MassWildlife, Angler Education Coordinator

Kristin McCarthy, MassWildlife, Education Assistant

Revised 2007

WATER SECTION OUTLINE

LEARNING OBJECTIVES

I. WATER BASICS

- DISTRIBUTION OF WORLD'S WATER
- STRUCTURE OF A WATER MOLECULE
- WATER CHEMISTRY
- HYDROLOGIC CYCLE

II. GROUNDWATER

- WATER TABLE
- POROSITY
- PERMEABILITY
- MOVEMENT OF GROUNDWATER IN THE SUBSURFACE
- RECHARGE AND DISCHARGE AREAS
- WELLS
- DETERMINING GROUNDWATER ELEVATION IN THE SUBSURFACE

III. SURFACE WATER

- WATERSHEDS
- STREAMS AND RIVERS
- FLOW
- DISCHARGE IN A RIVER OR STREAM
- ALTERATIONS TO WATERSHEDS
- EUTROPHICATION
- WETLANDS
- LAKES & PONDS
- LAKE TURNOVER & STRATIFICATION
- ESTUARIES
- OCEANS

IV. WATER POLLUTION

- EXAMPLES
- THE MOVEMENT OF CONTAMINANTS IN THE SUBSURFACE
- REMEDICATION

V. WATER QUALITY MANAGEMENT

- AT HOME
- IN SOCIETY
- DETERMINING WATER QUALITY
- BENTHIC MACROINVERTEBRATES

VI. REFERENCES & RESOURCES

VII. GLOSSARY

LEARNING OBJECTIVES


Team members will be able to:

- Identify the processes and phases of the water cycle.
- Describe the chemical and physical properties of water and explain their implications for freshwater and saltwater ecosystems.
- Analyze the interaction of competing uses of water for water supply, hydropower, navigation, wildlife, recreation, irrigation, industry and others.
- Discuss methods of conserving water and reducing point and non-point source pollution.
- Identify common aquatic organisms through the use of a key.
- Define a watershed and delineate a watershed boundary.
- Briefly describe the benefits of wetlands, including both function and value.
- Describe the benefits of riparian areas, including both function and value.
- Describe seasonal changes to a variety of aquatic ecosystems.
- Know methods used to assess aquatic environments and to mitigate invasive species negatively impacting those environments.
- Understand the various water quality parameters and use the water quality information to assess the general water quality of a body of water.

I. WATER BASICS

DISTRIBUTION OF THE WORLD'S WATER


Two-thirds of the planet Earth is covered by water. Most of that water is found in the oceans. Less than 1% is groundwater, surface moisture and soil moisture.



World Resources Institute/WHO/UNPD

STRUCTURE OF A WATER MOLECULE

A water molecule consists of one oxygen atom and two hydrogen atoms that are covalently bonded. Covalently bonded means that each hydrogen atom shares one outer shell electron with the oxygen atom and the oxygen atom shares an outer shell electron with each hydrogen atom. Covalent bonds only occur between two elements that are non-metals.



Water Molecule

Water has one of the highest heat capacities. Water can absorb a lot of heat energy before the temperature of the water will rise. A lot of heat energy is needed to break the hydrogen bonds that water molecules form with each other. Each water molecule is capable of forming four hydrogen bonds with adjacent molecules. This property is extremely important because water on the surface of the earth absorbs a lot of heat from the sun, which regulates the earth's climate.

Water has a strong surface tension. The water molecules at the surface of the water experience hydrogen bonds with adjacent molecules and molecules beneath them. The strong attractions experienced by the surface molecules enable them to withstand some pressure. Strong surface tension is also what enables bugs to skim across the surface of ponds. Due to water's unique molecular makeup, it has the ability to attract many different substances. It is known as the universal solvent because it can dissolve so many substances. The most common example of this is salt, since 97% of the earth's water is in fact salt water.

WATER CHEMISTRY

The chemistry of water will vary throughout the world. As water comes into contact with naturally occurring inorganic* substances, some of these substances will dissolve into the water. The types of dissolved substances and their concentration is dependent upon:

- the chemical composition of the precipitation* in that area
- the biological and chemical reactions occurring in that area
- the mineral composition of the geological material through which the water moves
- the length of time that the water spends in contact with the geological material through which it moves (Heath, 1998)

Water that spends a greater amount of time in contact with geologic materials will have a great concentration of dissolved inorganic substances. Water that spends only a short time in contact with geologic materials will have less dissolved inorganic substances. The concentrations of substances dissolved in water are referred to as the total amount of dissolved solids. The quality of water depends upon the total amount of dissolved solids and how these affect the characteristics of the water.

THE HYDROLOGIC CYCLE

The hydrologic cycle is the constant movement of water above, on and through the earth's surface. The diagram below shows the different paths that water may follow once it falls to the land in the form of precipitation. Water is returned to the atmosphere by evaporation*. Water may also be returned to the atmosphere by transpiration. Transpiration* is the loss of water vapor from the stomata (gas-exchange pores) of plants and to a lesser extent, from the evaporation of water through plant cell tissue. The cycle starts over again when the evaporated water molecules fall to the earth in the form of precipitation.




Magnificent Ground Water Connection, 1998)

II. GROUNDWATER

Groundwater is water that is located in the subsurface of the earth where all pore spaces or openings are saturated or completely filled with water. Pore spaces are the spaces in between soil and rock particles. To visualize this think of a jar of marbles. The marbles represent the soil or rock particles and the spaces in between them represent the pore spaces. Openings can also be fractures in a rock layer.

WATER TABLE

The water table is the highest level of the groundwater. The level or height to which the water may rise or fall depends on how much water is present. Water tables are higher during rainy months and when there is less transpiration, and lower during drier months and when there is more transpiration. Above the water table the ground may contain some water, but the pore spaces or openings also contain air. This is known as the vadose zone or the unsaturated zone. Below the water table is the saturated zone. In the saturated zone all pore spaces or openings are completely filled with water. Directly above the saturated zone is the capillary fringe. The capillary fringe* contains a little water that is held in place due to dipole-dipole force of attraction between water molecules and between water molecules and other substances. When the water table intersects the surface of the earth, streams, lakes, springs, or swamps form.



USGS Water Science for Schools <http://ga.water.usgs.gov/edu/index.html>

AQUIFERS

Any geological structure that can yield a usable quantity of water is called an aquifer*. Examples of a geologic structure include a subsurface layer of sand, gravel, clay, till (a combination of sand, silt, clay and gravel), limestone, slate, shale, sandstone, or bedrock. In an aquifer water may be located in the pore spaces (the space in between the particles) or in the fractures of rock. If the water table is free to rise and fall in the geologic formation the aquifer is said to be an unconfined* aquifer. If an impermeable geologic formation, such as a clay layer, overlies an aquifer the aquifer is said to be a confined* or an artesian* aquifer.

POROSITY

Porosity is a ratio of the total openings to the volume of a geologic formation. This ratio is then expressed as a percentage. The percentage is the maximum amount of water that a particular geologic structure can contain in relationship to its total volume. For example, suppose that a section of sandstone aquifer had a porosity of 33%; meaning that when the aquifer was saturated, 33% of the total volume would contain water.

PERMEABILITY

The ability of the material to transmit a fluid is called its permeability. If a material allows water to flow through it easily, it has a high permeability. If a material does not allow water to flow through it easily, it has a low permeability. Permeability depends upon two factors: the size of the pore spaces in the material, and how well the pore spaces are connected. All porous materials are not permeable, because if the pore spaces in the material are not large enough for water molecules to pass through than no water can be transmitted through the material.

MOVEMENT OF GROUNDWATER IN THE SUBSURFACE


Water enters the subsurface by infiltration*. Once in the saturated zone water moves at a very slow rate, which can be measured in cm/day or meters/year. Groundwater flow is laminar*, meaning the water has a smooth, not turbulent* flow that allows very little mixing of adjacent layers. The height of the water table and/or the amount of pressure that is exerted on the groundwater by the overlying subsurface is the primary influence on the flow of groundwater.

Groundwater is also influenced by gravity. Groundwater will move from areas where the water table is high to areas where the water table is low. The water table is low in areas where it intersects the land surface such as lakes, ponds, springs, streams, or swamps. Some of the water molecules will travel the shortest path to the waterbody. Other water molecules will travel deeper into the subsurface, taking a longer path to the surface water body. Water will also move from areas of high pressure to areas of low pressure. An example would be the land area above an aquifer, where pressure is exerted on that aquifer by that land directly above it. This will cause the groundwater within the aquifer to seek out areas of lower pressure.

RECHARGE AND DISCHARGE AREAS

A recharge area is a section of land where groundwater is able to infiltrate the subsurface and enter the saturated zone* of an aquifer. In general, an unconfined aquifer has a greater recharge area than a confined aquifer since its saturated zone is not overlain by a confining layer such as the case in a confined aquifer. However some lakes, ponds, streams and rivers are also recharge areas. This means that water is able to infiltrate the subsurface and enter the saturated zone via the ground under the surface waterbody. A discharge area is where groundwater leaves the saturated zone of an aquifer and flows into an area of land or a surface water body such as a lake, pond, spring, stream, or river.


Some surface water bodies alternate between being a recharge area or a discharge area depending on the state of the water table. For example, if the water table is higher than the stream bed the stream will serve as an area of discharge for groundwater resulting in the stream gaining water. If the water table is lower than the stream bed, the stream serves as an area of recharge to the groundwater resulting in the stream losing water.



Dauphin County (PA) Conservation District

WELLS

A well is a vertical hole drilled in the earth to access water. Wells vary in length and diameter. Some wells, water supply wells, are pumped to bring the water to the surface. Other wells, observation or monitoring wells, are used for taking groundwater samples and measuring the depth to groundwater. When a well is pumped it will reduce the level of the groundwater. This is called drawdown. Drawdown creates a cone of depression around the pumping well that extends in all directions. Drawdown may even result in a shallow well becoming dry if a deeper well in the area has lowered the level of the water table.



USGS Water Science for Schools <http://ga.water.usgs.gov/edu/index.htm>

DETERMINING GROUND WATER ELEVATION IN THE SUBSURFACE


It is important to know the elevation of groundwater in the subsurface, as this can aid in the drilling of water supply wells and in construction. Constructing a building where the water table rises close to the surface can cause problems, especially during the wet season where the water table can end up in your basement. Groundwater elevations are also used to determine the direction of groundwater flow. This is especially important if the water is contaminated*. If the flow direction of contaminated water is known, warnings can be issued and a treatment designed.

III. SURFACE WATER

Water that occurs on the surface of the earth is termed surface water. Surface water includes rivers, streams, wetlands, lakes, ponds, reservoirs, estuaries and oceans.

WATERSHEDS

The land contains water both on the surface and under the surface, and gravity forces all of this water to a main course (stream/river), or holding area (lake/pond/reservoir). A watershed consists of all the land that drains to a particular waterbody. It's the land that water flows across or under on its way to a river, lake, or reservoir.




Every watershed has boundaries which are defined by the highest elevations around a given waterbody. Ridges and hills serve as watershed boundaries or divides. Surface water will always flow down hill. Water that falls on the top of a ridge or a hill may flow down either side. Water that flows down one side will enter into one watershed while water flowing down the opposite side will enter into a different watershed. Thus, the high points in elevation form watershed boundaries by separating the flow of water into two or more watersheds. One of the most well known divides is the Continental Divide, running north to south through North America. Water that falls west of the Continental Divide flows to the Pacific Ocean. Water that falls east of the Continental Divide flows to the Atlantic Ocean.

STREAMS AND RIVERS

Streams and rivers within a watershed are classified by orders. The orders are defined below:

- **First-order stream** – known as headwaters (the beginning of the stream, the uppermost reaches of the stream); no other tributaries are connected to it.
- **Second-order stream** - the resulting stream from a first-order stream joining with another first-order stream.
- **Third-order stream** - the resulting stream from a second-order stream joining with another second-order stream.
- Orders increase (four, five, and beyond) with the increasing size of the stream or river.



Marsh 1998; www.chesapeake.towson.edu

Streams form different patterns. Five patterns are shown below. Note that when looking at a map of a dendritic stream pattern you can figure the direction of stream flow. Look for a "V" on the map. This is the joining of two streams. The bottom of the "V" points in the direction of stream flow.



Dendritic



Trellis



Radial



Braided




Meandering

Firehock, 1995

Streams can be further classified as perennial or intermittent. A perennial stream flows all year long as opposed to an intermittent stream that only flows when the water table is high. Intermittent streams can be found in later spring, early summer and after heavy rains.

FLOW

The flow in a river/stream can be turbulent or laminar. Flow depends on the velocity of the water and on the nature of the stream bed. Turbulent flow is when the water moves erratically downstream and stirs the sediment as it moves. Turbulent flow usually occurs when the stream bed is rocky. This type of flow can cause scour. Scour is the erosion of the channel by the suspended sediments and particles in the turbulent flow. Laminar flow is when the water moves steadily and there is minimal mixing of the sediment. The bottom is usually smooth and flat for laminar flow to occur.




Hewitt et al 1994; Redrawn by Jim Lafley, MA DCR

DISCHARGE IN A RIVER OR STREAM

The discharge is the volume of water that flows past a given point in a channel per a given unit of time. The velocity of a stream is dependent upon the stream's gradient. The gradient of a stream is the ratio of the vertical drop to the horizontal distance for that drop. The greater the gradient, the greater is the velocity. The size and shape of the channel also affect the velocity. The greater the contact area of the water with the land, the greater is the friction and the slower the velocity. The maximum velocity of the flowing water in a stream depends upon the shape of the stream channel. The maximum velocity of the flowing water in a straight channel stream is in the middle of the channel and close to the surface. The maximum velocity in a stream channel that has bends in it is toward the outside of each bend and slightly below the surface of the water. Erosion* occurs where the stream velocity is greatest and deposition* occurs where the stream velocity is slowest.

Erosion of cut bank



Hewitt et al. 1994; Redrawn by Jim Lafley, MA DCR

ALTERATIONS TO WATERSHEDS

As the land becomes more and more developed, diking, damming, straightening, and underground piping are altering waterways within our watersheds. Dams are used for many different purposes. They provide hydro-electrical power, recreation, irrigation, or traps for sediment from agriculture or construction activities. Although dams are beneficial in some situations, they are detrimental in others. While the water is behind the dam, it is being heated by the sun. A three degree rise in Celsius can cause thermal pollution and depletion of oxygen. Warmer water can not hold as much oxygen. Without enough oxygen to meet the biological oxygen demand the plant and animal life become stressed and could die. Damming a river also restricts the flow of water downstream, resulting in a disruption to the balance of the ecosystem* (ex. the loss of shallow wading habitats).

When the natural vegetation in the riparian zone* is removed erosion can take place. The roots of the vegetation help to hold the soils in place. Without the vegetation holding the earth in place, the surface water run-off can become high in suspended sediments. Extra sediment can cloud the water reducing the amount of photosynthesis by aquatic plants, clog the gills of aquatic animals, and smother macroinvertebrates*. If the sediments increase to the point where the river becomes too shallow for boat traffic, dredging must take place. Dredging is a solution for deepening the river but it also disrupts the established ecosystem and suspends the sediment.

Proper sediment management during agricultural activities and construction can save the streams. Silt fences and straw bales can help to contain sediments in construction areas. Alternating crops with rows of grass will also help to filter the runoff and reduce the amount of sedimentation in the water. Preserving vegetation along stream beds and especially in the riparian zones will also prevent extra sediments from entering the water.

When streams are in the way of development they are piped underground. Once underground they are often connected to the city's storm water drainage system. Underground streams in cities have been redesigned to carry away all the storm water, and whatever the storm water can carry, such as oil and animal wastes. In some cities, the

storm water drainage system is also used as an overflow for the city's sewage. Companies have been known to illegally connect to the storm water drainage system to discharge manufacturing by-products.

EUTROPHICATION

Eutrophication is the evolution, or natural aging process of a waterbody. A lake or pond is created from a catastrophic event, typically glaciers, volcanoes, or earthquakes, then slowly evolves from a pristine low nutrient waterbody to dry land over time. This usually takes thousands of years to occur, although human activities can speed the process up significantly. The process begins with nutrients, like nitrates and phosphorous, slowly building up in the waterbody. Plants and algae utilizing these nutrients grow, providing a good environment for aquatic life. Aerobic * (oxygen consuming) bacteria break down the organic wastes from this aquatic life. The decaying organic material and respiring algae cause the oxygen levels to drop. This will eventually reach a point where the waterbody no longer has enough oxygen to support life, and aquatic animals like fish begin to die. Without oxygen aerobic bacteria will no longer be able to break down the organic wastes, so the nutrients will stay trapped in these wastes within the waterbody. Anaerobic bacteria* (non-oxygen consuming) begin to break down the organic waste very slowly over time producing foul smelling gases. After hundreds of years, the waterbody will fill up with plants, becoming a wetland, then eventually dry land, most likely a forest.

As stated above, humans can greatly increase the process of eutrophication, this is sometimes known as cultural eutrophication. Sewage, agricultural runoff, and over-fertilization can severely impact both our salt and fresh water ecosystems, and cause an increase in plankton and encourage the growth of aquatic plants. The effects are most obvious in bays, coves, and estuaries as they are often shallow and sheltered.

WETLANDS

A wetland is defined as land that is saturated by water for at least part of the year, with characteristic hydric (saturated) soils, and having one of a number of distinct vegetation types (Art 1993). It was once common practice to drain and fill wetlands. It is now known that wetlands are much more valuable in their natural state. Wetlands are able to degrade moderate amounts of contamination (rendering these substances harmless), and trap sediments, preventing them from entering a stream. Wetlands can be a recharge or a discharge area for an aquifer, or alternate between the two. Wetland vegetation can store and slow the release of water, which helps to control flooding. Wetlands provide food and shelter for 45% of endangered animals and 26% of endangered plants (Gralla, 1994).

Swamps*, bogs*, marshes*, fens*, and vernal pools* are all types of wetlands. When they formed, the water table was at or near the surface of the earth. Swamps remain wet for part of the year and dry for the other part. The trees in a swamp are often dormant. Swamps are characterized by certain types of trees such as: oaks, willows, alders, tupelo gums and bald cypresses. Bogs have spongy, moist ground with only shallow waters visible. Sphagnum moss can usually be found in a bog. Marshes are wet the entire year. Grasses will grow in a marsh but trees will not. Cattails, bulrushes, and water lilies may be found in a marsh. Vernal pools are common in shallow lowland areas following spring runoff or high precipitation.

LAKES & PONDS

Lakes and ponds are classified as lentic* (still water) systems, in contrast to streams and rivers which are lotic* (flowing water) systems. There is no clear definition between a lake and a pond. There are several interpretations over which the science community likes to argue. Generally a lake is a much larger system, loosely defined as a body of fresh water that is deep enough to prohibit rooted aquatic plants from growing all the way across the bottom (Art 1993). A pond is a smaller shallow system that can potentially support rooted aquatic plants throughout. Again these are not hard fast definitions, but should help you to understand that a lake and a pond are different systems. Some of the ways that lakes and

ponds have formed include: glacial scouring from ice events (this is how most waterbodies in the northeast were formed), shifting of the earth's surface to create basins, and rivers changing course over time leaving behind ox-bow ponds, or lakes.

Lakes are divided into successional, or developmental stages. This has to do with a lakes trophic level (how nutrient rich or poor it is), which has a lot to do with how old the lake is, how much sediment has settled into the lake, and how much organic and inorganic matter the lake has been exposed to. This successional process is known as ontogeny*.


There are two major developmental stages of a lake, these are:

- **Oligotrophic** – A lake that is fairly young developmentally. These water bodies are extremely nutrient poor, and contain little organic matter. Oligotrophic lakes have a much clearer appearance than eutrophic lakes, and the bottom is comprised mainly of rock and sand. Most New England lakes are far removed from this oligotrophic stage, as these waterbodies were created during the last glacial event roughly 12,000 years ago. Reservoirs (man-made waterbodies) are considered oligotrophic as most were created within the past 100 years.
- **Eutrophic** – A lake which is rich in organic matter and vegetation. The water is much less clear than an oligotrophic lake, and is often considered murky. A eutrophic lake is much more evolved than an oligotrophic lake due in large part to an increased sediment load, and a higher nutrient (organic and inorganic) content. Eutrophic lakes are always more productive biologically, which simply means an increase in aquatic life. Most New England waterbodies are either eutrophic, or well on their way.

Lake and pond succession is a natural event, where nutrients and sediments from the surrounding land wash into and accumulate in the waterbody. An oligotrophic lake will eventually turn into a eutrophic lake. As a eutrophic lake accumulates nutrients and sediments it will eventually evolve into a wetland. These successional processes evolve over extremely long periods of time, often thousands of years.

A lake can be divided into six different zones:

- **Littoral zone** - The area of the lake that is closest to shore. The water here is usually shallow and vegetation will grow in this area.
- **Pelagic zone** - the open water in the middle of the lake
- **Limnetic zone** - the depth to which sunlight can penetrate the water of the lake
- **Photic zone**- Sunlit upper waters extending from the surface down to where light dims to about 1% of that at the surface.
- **Profundal zone** - the deep water below the level of light penetration
- **Benthic zone** - The bottom of the lake. This is where anaerobic microorganisms live in the silt and sediments. They break down the decaying vegetation and organic matter in the lake.



Jepson: <http://www.ent.orst.edu/jepsonp/2000lecture1>


LAKE TURNOVER & STRATIFICATION

Temperate lakes go through two periods of thermal mixing each year, one in the spring, and one in the fall. This thermal mixing occurs when the entire water column becomes one temperature. This process is also known as turnover. If a lake turns over twice a year it is known as dimictic. In other lakes around the world this turnover occurs more or less frequently, depending upon elevation, differences in climate, and lake morphometry*. To understand how lakes change it is important to know what happens to water at different temperatures. Water, like most substances, becomes lighter when warmed, and heavier when cooled. Water does have a unique property however, when it cools below 39 degrees F (4 degrees C), it becomes lighter. This is because water reaches maximum density at 39 degrees F. This results in the lake's bottom water being warmer than the surface water during winter. It also explains why ice, at 32 degrees F (0 degrees C), being less dense than water, floats.

Following these turnover events a lake begins to set up distinct temperature zones. These zones get larger and more distinct with prolonged periods of warm (summer) or cold (winter) weather. An upper warm water layer called the epilimnion, a cooler middle layer called the metalimnion, and a deeper colder water layer called the hypolimnion characterizes summer stratification*. A thermocline, or rapid drop in water temperature with respect to depth defines the metalimnion. If you have ever jumped into a lake in the middle of summer you probably have pushed through this layer (thermocline) and experienced this rapid drop in temperature. During the winter months these layers of stratification will again set up, but in reverse. As the temperature of the water reaches the point of maximum density surface ice will begin to form. Inverse stratification then occurs, where colder less dense water lies over warmer more dense water. During this time water temperatures throughout the depths of the water column will only be a few degrees different. They will stay this way until the spring turnover period.


Turnover, whether in spring or fall, is triggered by air temperature. Fall turnover is triggered by a decrease in air temperature, while spring turnover is triggered by an increase in air temperature. Wind also helps to churn the water column, and plays a big role in both seasons. Basically air temperature and wind act to break up stratification. In the fall, cooler air temps and gusty winds cool the upper layers, and eventually mix the entire water column to one temperature. In the spring, the warming air temperatures and winds melt and break up the icy top layer, and again, eventually mix the entire water column to one temperature.

Turnover is extremely beneficial to the lake because the lower layer needs to mix with the upper layers to become reoxygenated and release accumulated noxious gases. The lower water layer often has lost all of its oxygen due to the lack of contact it has had with the atmosphere during stratification. A build up of methane and hydrogen sulfide can also occur during stratification. These noxious gases have accumulated as by-products during the decay of organic matter at the bottom of the pond or lake. If turnover occurs early and rapidly (a few hours) due to a severe storm and high winds, the lower layer will not have had sufficient time to reoxygenate. This means that a small volume of oxygenated water will mix with a large volume of deoxygenated water, resulting in an insufficient amount of oxygen to sustain animal life.



Ohio State University

Layers in a Stratified Lake



Designed by the Washington State Lake Book -Recreated by Michelle Robinson

ESTUARIES

An estuary is where fresh water from rivers mixes with salt water from oceans. Estuaries contain a great diversity and number of life forms because they are rich in nutrients brought by the flow of fresh water from rivers. Some animals that live in estuaries can tolerate both fresh and salt water. Bays, mudflats and salt marshes are all examples of estuaries. Estuaries are tidal, meaning that a part of the land under the water may be exposed to air at low tide.


Due to the beauty and recreational activity estuaries provide, people like to develop the land that surrounds it. Currently, approximately 60% of the world's population lives near the coast (Gralla, 1994). Unfortunately the closer people live to the water the easier it is for pollution from lawn products, industry, sewage, and storm water drainage systems to come in contact with the water. Shellfish and other filter feeders ingest different kinds of contamination and bioaccumulate* it in their systems. This poisons both the animals, and the people who eat these animals.

OCEANS

Oceans cover two-thirds of the world. Oceans gave birth to the first living organisms over 4 billion years ago. Oceans have an average depth of 2.3 miles (3.7 km) but can be as deep as 7.14 miles (11.5 km) (Towel, 1991). Some of the largest mountain ranges in the world are found in the oceans. These ranges are called Midocean Ridges. The salinity content of the oceans is approximately 3.5% (Towel, 1991). Most of the salt is sodium chloride (NaCl).

The ocean can be divided into seven zones:

- **Pelagic zone** - the open ocean.
- **Neritic zone** - The part of the pelagic zone that extends over the continental shelf. The continental shelf is a flat plane or terrace that borders the land and slopes down into the ocean. The size of the continental shelf varies. The seaward limit of the shelf is defined by a dramatic change in the slope of the land. This zone supports the largest amount of marine life due to its nutrient rich waters from the accumulation of sediment and organic debris from the land and the upwelling of nutrients from the ocean bottom.
- **Oceanic zone** - the part of the pelagic zone that is not over the continental shelf and is defined as the deep water of the open ocean
- **Photic zone** - The area of the ocean that sunlight is capable of penetrating. This zone extends to a depth of approximately 656 feet (200 meters) from the surface of the water.
- **Aphotic zone** - the area of the ocean that receives no sunlight
- **Benthic zone** - the bottom of the ocean
- **Intertidal zone** - the area of land where the tide rises and falls



Jepson: <http://www.ent.orst.edu/jepsonp/2000lecture>

Temperature, salinity, hydrostatic pressure (the pressure exerted by a static column of water), light, currents and nutrients affect the ecosystem of the ocean. The base of the food web is phytoplankton*. Through photosynthesis and the ingestion of nutrients, phytoplankton are able to reproduce in great numbers. Zooplankton* feed upon phytoplankton. The term zooplankton refers to a variety of life forms such as krill (tiny shrimp-like organisms) and larvae. Zooplankton are consumed by small surface feeding fish such as anchovies and sardines. The small fish are in turn fed upon by larger fish and marine mammals. Scavengers such as crabs feed on the dead plants and animals on the bottom. Micro-organisms* degrade the remains left by the scavengers, producing nutrients, which are brought up from the bottom during periods of upwelling.

The only ecosystem that does not rely on the sun for survival is located in the ocean. At the bottom of the open ocean are hydrothermal vents, which emit a continuous fountain of hydrogen sulfide gas. Hydrogen sulfide gas is poisonous to humans. Bacteria in the area of the vent are capable of converting this gas into energy, which allows them to reproduce and form the base of the food chain. The creatures that the bacteria support at the hydrothermal vent are extremely different. Giant clams, albino blind crabs and worms 4 meters in length have been found.

The oceans provide us with a source of precipitation due to the large amount of evaporation of water. Oceans also help to regulate our climate due to water's high heat capacity. One-half to one third of our oxygen comes from phytoplankton in the ocean. These one celled plants take in carbon dioxide and produce oxygen that humans and animals need to survive. This process occurs during photosynthesis. Another benefit to this process is the regulation of carbon dioxide, a known greenhouse gas.

Humans have not been entirely kind to our oceans, pollution from land is carried to the sea via rivers, and sometimes the oceans are used as a receptacle for our garbage. Over-harvesting certain species of marine wildlife has upset the ecological balance, at times pushing other species to near extinction.

IV. WATER POLLUTION


EXAMPLES

There are numerous ways that pollution can enter water supplies. Below are a few;

- Industry can discharge waste by-products directly into waterways. Many industries have permits to discharge a specific quantity of by-product but exceed their limit and pollute the water. Illegal or accidental discharge of chemicals into the ground can also contaminate the water and subsurface. Both are examples of point source pollution. Point source pollution means that the exact geographical location of the source of contamination can be located. Non-point source pollution means that the exact geographical location of the contamination can not be located. An example of a non-point source pollutant is agricultural erosion that can't be identified like a discharge pipe which releases a specific contaminant.
- Some power plants discharge hot water into a water body, changing the natural temperature of the water, and upsetting the established ecosystem. This is known as thermal pollution.
- Chemical storage tanks or underground storage tanks are known to leak. Chemicals may leak from tanks located on the surface or in the subsurface and contaminate the environment. This contamination can flow with the surface water run-off and enter rivers or other surface waterbodies, and/or infiltrate the subsurface and enter the groundwater just as water does.
- Out-dated septic systems may leak, or animal waste may be washed overland into the water, allowing disease-carrying bacteria, called pathogens, to enter the water supplies. Boaters who empty their sewage into the water instead of using an approved pump out station at a marina also contribute to this problem.
- Pesticides, fertilizers, and other agricultural chemicals can enter groundwater by infiltration, or can be washed off crops and flow with the surface water run-off entering rivers or other surface waterbodies.
- Leachate* from landfills or garbage dumps may carry contaminants to water supplies.

- Phosphates and nitrogen can cause an excess growth of bacteria in lakes or ponds. These bacteria use all the available oxygen in the water, leaving the water with an insufficient amount of oxygen to support its ecosystem.
- Oil tankers may run aground or have leaking hulls, allowing large quantities of oil to spill in the oceans.
- At times the ocean is used as a dump for garbage or a place to pump treated sewage.
- Motor boats and jet skis pollute by adding oil and gasoline by-products to the water.
- Polychlorinated bi-phenyls (PCBs) from the production of electrical transformers have entered rivers and bays. The PCBs adsorb to the sediments in the water. Once the fish ingest the contaminated sediments, the PCBs bioaccumulate in their tissues. Since PCBs are known carcinogens* the fish become toxic and can not be eaten.
- Mining produces wastes that are harmful to the environment. An example of this is acid mine drainage. Run-off at mining sites turns acidic due to the oxidation of pyrite and iron disulfide by exposure to the air and water. The oxidation process produces ferrous ions and sulfuric acid, which enters the surface water run-off. Acid mine drainage can also infiltrate the land and pollute the groundwater.
- Acid precipitation is formed when the gases, sulfur dioxide and nitric oxide (produced by the burning of fossil fuels) mix with water in the atmosphere to form rain, snow, fog, frost, sleet, and cloud moisture with a pH of 2.0 or less. Normal uncontaminated rain has a pH of about 5.6. This precipitation erodes buildings, and leaches heavy metals out of the soils, which can end up in surface water. Depending on how acidic the water, it can be toxic to animal life. If the pH levels are low enough fish and amphibians can not reproduce, and immature fish (fry/larvae) can die.
- In coastal areas, if too much water is withdrawn from a water supply well that is close to the ocean, saltwater intrusion* can occur. This occurs when saltwater is drawn into the well by pumping. It is estimated that two thirds of all aquifers in the United States are underlain by salt water (Winter et al., 1998). This is especially a problem in coastal areas where fresh rainwater infiltrating the ground

floats on top of salt water in the aquifer. Salt and freshwater do not mix (differing densities) unless too much fresh water is being withdrawn from the well.



Ground and Surface Water Contamination Sources
Dauphin County (PA) Conservation District Dauphin County Conservation District

THE MOVEMENT OF CONTAMINATION IN THE SUBSURFACE

Once contaminants infiltrate the subsurface and enter the groundwater, it is important that hydrologists* be able to predict the direction of movement of the contamination. To predict this, the density of the contamination relative to the groundwater must be known.

REMEDICATION

Remediation is the process of removing contamination from the environment. Passive remediation or active remediation may remove contamination. Passive remediation refers to the adsorption, dilution, and dispersion of the contaminant by the environment. These processes do not rid the environment of contamination, only lessen the concentration of the contaminant by spreading it throughout the environment. Another form of passive remediation is bioremediation. This occurs when naturally occurring bacteria in the ground break apart the molecules of the contamination. The bacteria derive energy by breaking down the molecules. This can rid the environment of toxic substances because the bacteria can eventually change a toxic contaminant into to a non-toxic substance. Active remediation refers to any activities conducted by people to clean up the contamination. Forms of active remediation include:

- Using a trench to catch and divert contaminated water.
- Installing an underground wall or barrier to stop the flow of contaminated water.
- Pumping groundwater to the surface by using a well and either, treating the water and returning it to the subsurface, or collecting the water and removing it from the subsurface. Forms of treatment may include the addition of chemicals to cause the contamination to precipitate out of the groundwater, filtration of the water, and/or aeration* to volatilize* contaminants out of solution. With aeration care must be taken to capture and treat the volatiles.
- The physical removal of contaminated water and soil is done by absorbing the contamination with booms or collecting it in 50-gallon drums. This option is used when an oil spill occurs in the ocean.

V. WATER QUALITY MANAGEMENT

Every life form on the planet depends on water for its survival. When the quality / or quantity is compromised, these life forms have problems surviving. Maintaining water quality and quantity, therefore, is essential to maintaining life on earth. Almost everything we do affects water quality. Our individual use of water at home and our use as a society can have a significant impact.

AT HOME

- If we use water wastefully, we are contributing to the depletion of the region's water supply, whether you get your water from a private well, or a municipal supply. Sound water conservation practices go hand in hand with sound water quality programs.
- Lawn and garden chemicals (fertilizers/herbicides/pesticides) can run off into storm drains and local waterways during storms.
- Household hazardous wastes such as solvents, paints, waste oils, and cleaning chemicals flushed down the drain or poured into storm drains can make their way into either surface water or groundwater.
- Improperly maintained or overused septic systems or cesspools allow excess nitrogen and bacteria to enter regional water supplies. Sewer systems are designed to handle sewage (human and domestic waste products) not hazardous waste.

IN SOCIETY

- Larger scale lawn and garden chemical runoff from parks, golf courses, and crop lands, enters regional water supplies during storms, raising the level of nutrients, such as nitrogen or phosphates, altering the pH, or contributing toxic chemicals, such as pesticides, that might harm the environment.
- Each rainfall washes road debris, such as oils, bits of tire rubber, and gasoline into local waterways. This debris adds hydrocarbons and sediment to the water, which can often raise the turbidity levels.
- Industrial pollution threatens our waterways, adding a variety of contaminants, often in high concentrations, to regional water supplies.

DETERMINING WATER QUALITY

There are many factors that affect the water quality of a river, stream, lake, pond, or reservoir. Conditions can fluctuate over time making it necessary to measure water quality on a regular basis. Quality is also dependent on the water's specific use. Water may be safe for one use but completely unacceptable for another. Some examples of use categories are: water supply for domestic and industrial use, recreation for total body contact like swimming, partial body contact like fishing, hunting, trapping, and boating, protection of wildlife, agricultural uses such as livestock watering, irrigation, and spraying, and commercial uses like navigation, and hydroelectric power.

To determine the water quality of a given river, lake, pond, or reservoir, there are many tests to identify specific areas of concern. There are nine parameters* that make up what is called the water quality index (WQI)*. This index was set up to compare rivers throughout various parts of the country. The nine parameters are: temperature, pH, dissolved oxygen, biochemical oxygen demand, total coliform, total phosphorous, nitrates, turbidity, and total solids. After completing the nine tests a formula is used to determine the overall WQI #.

The overall WQI ranges are as follows:

- 90 – 100 Excellent
- 70 - 90 Good
- 50 – 70 Medium
- 25 - 50 Bad
- 0 – 25 Very Bad

A body of water could have an overall WQI in the good range, but one of the parameters, say fecal coliform, could exceed the state standards, which would then make the waterbody unusable for an activity such as swimming. So the overall score can be deceptive.

Here is a closer look at the parameters that make up the WQI:

Temperature – The temperature of a waterbody is very important. Physical, biological, and chemical characteristics are influenced by temperature. *Dissolved oxygen* levels are higher in colder waters, because gases, like oxygen, dissolve more readily at lower temps. Fish have temperature preferences, for example trout need colder water, while bass, and carp seek out warmer environs. Temperature also affects the rate of photosynthesis by aquatic plants, as well as the metabolic rates of other aquatic organisms.

pH - This test indicates the number of hydrogen ions present in a substance. A value of 7.0 is neutral, meaning there is an equal number of hydrogen (H⁺) ions, and hydroxyl* (OH⁻) ions. A number less than 7.0 indicates that there are more hydrogen ions than hydroxyl ions, making the substance acidic. If the number is greater than 7.0, there are more hydroxyl ions present and the substance is alkaline. The pH scale goes from 0.0 to 14.0. The scale is logarithmic, meaning each one-digit change in the scale indicates a ten-fold change in the acidity or alkalinity (example: a substance with a pH of 5.0 is 10 times more acidic than a substance with a pH of 6.0, and 100 times more acidic than a substance with a pH of 7.0).

There are many factors that influence pH, a few of these are:

- Decomposition of organic materials releases carbon dioxide. As carbon dioxide mixes with water, it forms carbonic acid, lowering the pH.
- Alkaline materials like limestone buffer acidic waters, unfortunately limestone is not present everywhere. It is scarce in the northeast.
- Fresh water is more susceptible to changes in pH than salt water as the minerals in saltwater act as buffering agents.
- Some human practices have an affect on pH values. Modern industry has contributed such noxious releases as sulfur dioxide, oxides of nitrogen, and carbon dioxide into the atmosphere. These compounds go through a series of changes in the atmosphere and eventually return to earth as acid precipitation. Acid precipitation is responsible for increasing acidity in many lakes of the northeast, where there is limited buffering capacity. Many of these lakes have a crystal clear, almost pristine appearance. Ironically, this look is due to an almost total lack of life. Fortunately over the past 20 years much has been done to

clean up these industrial practices, bringing back from the dead many of these water bodies.

Dissolved Oxygen (DO) – A measure of the amount of oxygen dissolved in water. Oxygen is essential to most all life, and water is capable of holding a large volume of oxygen. Water movement and aquatic plants are the two main ways oxygen enters a waterbody. There are minimum levels of oxygen necessary to support life. Most aquatic animals can withstand DO levels as low as 2.0 parts per million (ppm) for very short periods of time, but these levels are certainly not healthy. In general DO levels greater than 5.0 ppm is considered relatively healthy, supporting most aquatic animals. DO levels are influenced by salinity, temperature, and pollution; a higher percentage of oxygen can dissolve in water with lower temperature and low salinity.

Several factors affect the amount of oxygen in water. Some of these factors are:

- **Salinity** – the less of this, the more DO
- **Agitation and Turbulence** – the more contact with atmospheric oxygen the higher the DO
- **Temperature** – the lower the temperature the higher the DO
- **Minerals** – the higher the mineral content the lower the DO
- **Plant Life** – Photosynthesis (green plants utilizing sunlight, chlorophyll, water, and carbon dioxide to produce water, complex carbohydrates, and oxygen) occurs during the daylight, so DO levels can be quite a bit higher during this time than at night. So more photosynthesis equals higher DO.
- **Organic Wastes** – Certain aerobic bacteria that decompose organic wastes (feces, detritus) consume oxygen, so more wastes equals lower DO.

Biochemical Oxygen Demand (BOD) – Basically this is a measure of the oxygen-consuming life in the water. Organic materials (leaves/sewage/dead animals/etc..) are decomposed by bacteria that live in the water. The amount of oxygen consumed by the bacteria over a certain period of time in a controlled environment indicates the amount of this organic matter in the water. That measure (amount of oxygen) is called the biochemical oxygen demand (BOD). A high BOD might indicate the presence of pollution in the waterbody, whereas a low BOD can be an indicator of clean water. BOD serves as an important clue as to a water bodies health.

Interpreting BOD results:

- 1 – 2 ppm = very clean water w/little organic decay
- 3 – 5 ppm = moderately clean water, some organic decay (probably from plant life)
- 6 – 9 ppm = much organic decay (possibly from algae blooms)
- 10+ ppm = very unhealthy levels of organic decay (often from untreated sewage)

Nitrates – A common nutrient abundant in aquatic ecosystems, and essential for plant growth. Plants do not utilize nitrogen in its pure state, rather they use it in the form of nitrates, and ammonia. Bacteria initially break tissue down into ammonia, which is then oxidized by other bacteria into nitrites and nitrates, thus completing the cycle. Nitrate contributors include animal excrement, fertilizers, and sewage.

Total Phosphorous – Includes organic phosphorous and inorganic phosphates. Organic phosphorous is attached to organic matter composed of once-living plants and animals. Inorganic phosphates include ions bonded to soil particles, and phosphates present in laundry detergents. In most waterbodies phosphorous is the growth-limiting element, in other words, plant growth is limited by the phosphorous available within it's environment. Since very little phosphorous is necessary for aquatic plant and algae growth, excess phosphorous can thus produce extensive plant growth and algae blooms (pea-green colored water). This is the classic symptom of cultural eutrophication*. A few sources of phosphorous include: human and animal wastes, industrial pollution, wastewater treatment plants, residential septic systems, agricultural practices, as well as residential and commercial application of lawn fertilizers.

Total Solids – Includes both dissolved solids (portion of material in water that passes through a filter), and suspended solids (portion retained by a filter). Dissolved solids include a wide range of substances like calcium, sodium, phosphorous, iron, sulfate, carbonate, nitrates, and chloride to name a few. A constant level of these substances is essential for maintaining aquatic life. Suspended solids include silt and clay particles from soil runoff, plankton*, industrial waste, and sewage. High concentrations of total solids can diminish water quality and cause water balance problems for individual organisms. Low

concentrations may limit growth of aquatic life, and may also restrict some aquatic organisms from surviving in water.


Turbidity – The measure of a water's relative clarity, or cloudiness resulting from suspended solids in the water that reduce the transmission of light. These solids may be microscopic plankton, stirred up organic materials, eroded soil, clay, sand, industrial wastes, or sewage. High turbidity may be a symptom of other water quality concerns. A few effects of turbidity include, diffusing sunlight, which in turn reduces photosynthesis, raising water temperatures (due to the ability of the solids in the water to absorb and hold the sun's heat), clogging the gills of fish and other invertebrates, which stunts growth and decreases disease resistance, creating breeding grounds for pathogenic bacteria that live within the organic materials that cause higher turbidity, and industrial processes, where turbid water can clog machines and interfere with making food and beverages.

Total Coliform Bacteria – A group of generally non-pathogenic (non-disease causing) bacteria that live throughout the environment. One type of coliform that live in the intestinal tract of warm blooded animals to aid in digestion is known as fecal coliform. It's not harmful all by itself, but the presence of it serves as an indicator that other viruses or parasites may be present. Some of these others might however cause disease (pathogenic). In water with a high total coliform count, about 10% of the coliform will be fecal. Coliform bacteria are easy to test for, and usually occur with other pathogens. That is why people who monitor water quality use this group as an indicator. Sources of coliform bacteria include sewage treatment plant wastes, storm water, agricultural runoff, and improperly working septic systems.

BENTHIC MACROINVERTIBRATES

Beyond the chemical and physical testing of water to determine its quality, scientists can also use benthic macroinvertebrates. Macroinvertebrates are animals with no backbones that can be seen without the aid of a microscope. Benthic macroinvertebrates spend at least part of their lifecycle in or on the bottom of a body of water. They can include aquatic insects such as mayflies, stoneflies, caddis flies, midges and beetles, as well as worms and crustaceans such as crayfish, clams and snails.

The invertebrates are good indicators of water quality because they differ in their sensitivity to water pollution. They are not very mobile so they can not avoid pollution events or other forms of stress that some sampling methods might miss. If water quality is poor or if a pollution event occurred within the past several months, it will be reflected in the macroinvertebrate population. Some of these animals, such as stoneflies and mayflies, can't tolerate much pollution at all. If a sample is dominated by these organisms it would signify good water quality.



VI. REFERENCES & RESOURCES

Envirothon students are not required to study from the references and resources listed below.

- Art, Henry W., ed. *Dictionary of Ecology and Environmental Science*. New York: Henry Holt and Company, 1994.
- Cech, Thomas. *Principles of Water Resources*. New York: J. Wiley & Sons, 2003.
- Firehock, Karen. *Hands on Save Our Streams*. Gaithersburg, MD: Izaak Walton League of America, Inc., 1995.
- Gralla, Preston. *How the Environment Works*. Emeryville, CA: Ziff-Davis Press, 1994.
- Heath, Ralph C., "Basic Ground-Water Hydrology". U.S. Geological Survey Water Supply Paper 2220, 1998.
- Hewitt, Paul G., John Suchocki, and Leslie A. Hewitt. *Conceptual Physical Science*. New York: Harper Collins College Publishers, 1994.
- Murck, Barbara, Brian J. Skinner, and Stephen C. Porter. *Environmental Geology*. New York: John Wiley and Sons, Inc., 1996.
- Sims, J., R. Sims, R. Dupont, J. Matthews, and H. Russell. "In Situ Bioremediation of Contaminants in Unsaturated Subsurface Soils". EPA 540/S-93/501, 1993.
- Watson-Ferguson, Kami and Loren Kellogg, eds. *Aquatic Insects & Crustaceans*. Gaithersburg, MD: Izaak Walton League of America, Inc. 2006.
- Winter, Thomas C., Judson W. Harvey, O. Lehn Franke, and William M. Alley. "Ground Water and Surface Water: A Single Resource". U.S. Geological Survey Circular 1139: Denver, Colorado, 1998.

Resource Professionals

- Jim Lagacy, Angler Education Coordinator
MassWildlife, Field Headquarters, 1 Rabbit Hill Rd., Westboro, MA 01581
(508) 389-6309 Jim.Lagacy@state.ma.us
- Jim Lafley, Watershed Education Coordinator
DCR – Watershed Management, 180 Beaman St., W. Boylston, MA 01583
(508) 792-7423 x 231 Jim.Lafley@state.ma.us
- Jim Straub, Lakes and Ponds Program
DCR – Lakes and Ponds, 251 Causeway St., Suite 600, Boston, MA 02114
(617) 626-1411 Jim.Straub@state.ma.us
- Pat Doherty, MA Water Resources Authority School Programs
MWRA, 100 First Ave 39–1, Charlestown, MA 02129

Additional links can be found at <http://www.maenvirothon.org/aquatics.htm>

VII. GLOSSARY

Acid – a substance with a pH value below 7. It will be sour in taste or corrosive in nature.

Acid Deposition – wet and dry atmospheric fallout with a very low pH that results from the combination in the atmosphere of water vapor with sulfide or nitrous compounds released during the combustion of fossil fuels.

Aeration – expose water to the circulation of air for purification.

Aerobic – occurring, acting, or living in the presence of free oxygen, either as atmospheric gas or dissolved in water.

Alkaline – having a pH greater than 7, the opposite of acid. Alkaline soil or water is able to neutralize acid precipitation or discharge.

Anaerobic – occurring, acting or living in the absence of oxygen.

Aphotic Zone – the zone in a body of water where light cannot penetrate. Photosynthesis cannot be supported here.

Aquifer – a geologic structure composed of porous or permeable rock or sand that can yield a usable quantity of water.

Artesian Aquifer – see Confined Aquifer.

Bedrock – a general term for the solid rock that underlies the soil or unconsolidated surface material.

Benthic Zone – the section of a water body near the bottom.

Bioaccumulate – storing chemicals in an organism in higher concentrations than are normally found in the environment.

Biological Oxygen Demand – a measure of the dissolved oxygen consumed by micro-organisms as they break down organic matter in a sample of water. The greater the quantity of oxygen consumed the more organic matter present. It can indicate the presence of sewage or other organic pollution.

Bog – a wetland formed where low oxygen levels and soil temperature cause incomplete decomposition, limited drainage and an accumulation of fibrous peat.

Brackish – water containing a salinity level between freshwater and salt water.

Capillary Fringe – the zone above the water table (top of the saturated zone) in which water is held by surface tension

Carcinogens – substances that stimulate the formation of cancer.

Condensation – the process in which a substance changes from a gas to a liquid usually by cooling.

Confined Aquifer – a water saturated layer of soil or rock that is bounded above and below by impermeable layers causing a pressurized water source.

Contaminated – made impure by contact or mixture with a pollutant.

Current – movement of water caused by tides, wind, temperature or water flowing in a stream

Deposition – the process of laying down sediment or accumulating layers of material carried in suspension.

Dissolved Oxygen – oxygen in solution in water; oxygen available to gill-breathing organisms.

Ecosystem – a functional system comprised of a community of organisms and their environment, including the energy transfers and nutrient cycling.

Effluent – any matter that enters the environment from a specific source; the term generally refers to waste water from a sewage treatment or industrial plant.

Epilimnion – the zone of a thermally stratified water body occupied by the upper layer or warmer water.

Erosion – the removal or wearing away of soil or rock by water, wind or other natural forces.

Eutrophication – The natural aging process of a body of water. This process may be accelerated by human activity, including industrial or agricultural runoff, sewage, or the introduction of exotic plants.

Evaporation – changing of a substance from a liquid to a gas.

Facultative Species – a species that doesn't require one specific habitat type for its survival, but will seek out a given habitat type if it is within its home range.

Fen – low, flat, marshy land with herbaceous vegetation.

Groundwater – water found at or beneath the water table, in saturated soil or bedrock.

Headwater – the highest reaches of a stream in a watershed or drainage basin.

Hydrology – the branch of science that studies the distribution of the earth's waters both above and below the ground, including the water cycle.

Hydrologist – scientist who studies the properties, distribution and effect of water in the environment.

Hydroxyl ions – an OH ion found in bases, certain acids and atmospheric compounds.

Hypolimnion – the zone of a thermally stratified water body occupied by the deeper, cooler water layer.

Inorganic – involving neither organic life nor products of organic life.

Infiltration – the process of water or other liquid soaking into soil.

Intrusion – an unwelcome addition of saltwater into a freshwater source.

Laminar Flow – smooth, unidirectional flow in a stream or river.

Leachate – substance dissolved or washed out by a percolating liquid.

Lentic – meaning still or calm. In ecology it refers to still water environments (ponds & lakes).

Limnetic Zone – the zone in a lake or pond occupied by deep water where light cannot penetrate to the bottom. This area does not allow rooted aquatic vegetation to grow.

Limnology – the scientific study of freshwater aquatic ecology.

Littoral Zone – that zone in a lake or pond occupied by shallow water or shoreline. Sometimes referred to as the rooted aquatic plant zone.

Lotic – meaning moving or flowing. In ecology it refers to flowing water environments (streams & rivers).

Macroinvertebrates – animals without backbones that are large enough to be observed without the aid of magnification.

Marsh – a low wetland covered with water part of the year, consisting of grasses, rushes and sedges.

Micro-organisms – small organisms that can only be seen with the aid of a microscope. They are important in the degradation and decomposition of organic material from nature or artificial mechanisms.

Morphometry – is the measurement of the shape of a body of water.

Non-point Pollution – pollution that enters water through run-off from land and is difficult to pinpoint rather than from a specific place such as an effluent pipe.

Nymph – larval phase of an some aquatic insects.

Obligate Species – a species requiring a certain type of habitat for its survival, for example a Spotted Salamander requires a vernal pool for reproduction.

Ontogeny – the course of development of a lake or pond from water body to land.

Order – name given to a group of related families.

Organic – of, related to, or derived from living organisms. Organic substances contain carbon.

Parameter – a typical element which is tested for and used as an indicator or water quality.

pH – a measure of hydrogen ion concentration or acidity. The pH scale ranges from 0 (most acidic) to 14 (most basic) with a pH of 7 being neutral.

Photic Zone – the zone in a body of water where light penetrates. The light supports photosynthesis of organic matter in this zone.

Phytoplankton – tiny drifting organisms in aquatic environments that are capable of photosynthesis.

Plankton – microscopic organisms that are suspended in an aquatic habitat. (character on Sponge Bob)

Precipitation – water falling, in a liquid or solid state, from the atmosphere.

Riparian – located or living along or near a stream, river or body of water.

Runoff – water that drains or flows off the surface of the land.

Salinity – the amount of salt present in a solution.

Saturated Zone – the subsurface zone in which all openings are filled with water

Spring – a natural point of groundwater discharge formed where an aquifer intersects the surface.

Stratification – development of different temperature layers of water in a lake or pond.

Stream Gradient – the steepness of the stream bed or ratio of vertical drop to horizontal distance.

Sublimation – the transition of a substance from the solid phase directly to the vapor phase, or vice versa, without passing through an intermediate liquid phase.

Suspended Load – the amount of substance carried by a stream or river that is in suspension, also referred to as suspended sediment.

Swamp – a wetland where the soil is saturated and often inundated with water with trees as the dominant vegetation.

Thermal Stratification – orientation of the water by temperature in a lake or pond where less dense, warmer water positions over the top of more dense, cooler water to form a layering affect.

Thermocline – the zone in a thermally stratified water body where you observe the maximum decrease of temperature with respect to depth. You observe the temperature decreasing rapidly as you descend through the water column.

Total Dissolved Solids – a measure of the amount of total dissolved minerals in water.

Total Coliform Bacteria – a measure of the amount of coliform bacteria found in a given water sample. Coliform bacteria reside in the intestines of vertebrate animals including humans.

Transpiration – the process of giving off water vapor through plant leaves.

Tributaries – small feeder streams that flow into larger streams and rivers.

Turbidity – a measure of the ability of light to penetrate water, indicating the amount of living and non-living suspended solids and dissolved substances. It's the measure of a water's cloudiness.

Turbulent Flow – water that moves erratically downstream, stirring up everything it contacts.

Turnover – thermal mixing within a water body. The temperature and wind conditions during certain times of the year contribute to the overall mixing of the water column eliminating thermal stratification.

Unconfined Aquifer – an aquifer in which the upper boundary is the top of the water table.

Vernal Pool – a temporary pond existing during the wet spring season and drying up in the summer.

Volatilize – dispersing chemicals into the air through the aeration process.

Water Cycle – the continuous circulation of water in systems throughout the planet, involving runoff, evaporation, condensation, precipitation, transpiration and sublimation.

Water Quality Index (WQI) – an index based on nine parameters (dissolved oxygen, pH, nitrates, total dissolved solids, turbidity, phosphates, BOD, fecal coliform and temperature) used to measure the quality of a body of water, compare changes over time and compare to other water bodies.

Water Table – the uppermost level of groundwater.

Watershed – the total land area draining to a particular body of water.

Wetland – any land that tends to be regularly wet or flooded.

Zooplankton – plankton that is composed of tiny, often microscopic, animal and animal matter.

